

endocrine disorder) are discussed. Changes that occurred in the inwardly rectifying potassium channel gene (KCNJ11) were translated to the protein level by use of the web-based Translate tool of ExpASY Proteomics Server. A model of the protein was built based on the 3JYC.PDB structure, with the use of the HOMER server. A tetramer was subsequently built using the pre-calculated quaternary structure of 3JYC.PDB. Missense mutations that had been detected in the patients' samples, including E227K and E229K were then applied to the modeled structure and their effect studied with regard to altered interactions in and between monomers. Both mutations occur in a loop connecting two beta strands protruding from the cytoplasmic domain of the protein and are located in the interface between two monomers. Based on this model, R192, R134, and H193 of the adjacent monomer are found to have important roles in the interactions occurring in this region, which would be abolished upon mutation. Overall, this study is presented as a concrete example of protein modeling application in the study of endocrine genetic disorders. It should be mentioned that in this case, occurrence of missense mutations in the protein would be considered as one of the evidences used in the course of a change of medication in these patients.

Keywords: Protein Modeling, KCNJ11, Missense Mutation.

Abstract No.161

Study on Some Physicochemical Properties of Biomacromolecules and Their Application in Drug Targeting

*Mehrdad Mahkam¹, Solmaz Zakhireh*², Saeed Jafarirad³*

1. Azarbaijan University of Tarbiat Moallem, Tabriz, IR
 2. Islamic Azad University - Ahar Branch, Tabriz, IR
 3. University of Tabriz, Tabriz, IR
- (E-mail: zakhireh.solmaz@gmail.com)

To evaluate the release pattern of films with microspheres, the release pattern of glutaraldehyde cross-linked carriers (6%, 12% and 18% w/w) has been analyzed. The objective of this paper reply to this question that if degree of cross-linking and drug conjugation capacity could change the physicochemical aspects of biomacromolecular carriers as support to immobilize an enzyme model as well as in drug targeting. The study was conducted in two different stages. In the first stage the preparation of cross-linked chitosan films and microspheres (MGn, n=6, 12 and 18) successfully took place. In the second phase, the behavioral effects were investigated by using Differential Scanning Calorimetry (DSC), Scanning Electron Microscopy (SEM), UV-Vis spectrophotometry and other spectroscopic techniques. The reactive

amino groups of chitosan films and microspheres were participated in a chife-base reaction with glutaraldehyde that has at least two reactive functional groups. Some of these functional groups in the other side of the glutaraldehyde chain remain free to react with drug or enzyme's amine group (Scheme). Scheme . Synthetic method of network's macromolecule utilizing cross linking process conjugated by enzyme. Based on our results, degree of cross-linking and drug conjugation capacity, as two important factors, affect simultaneously the thermal, morphological and hydrolytic behavior of the carriers in film and microsphere forms. For example, the chitosan microspheres have low Tg (123 °C) relative to chitosan films because of its spherical structure (154 °C). The reducing of Tg in chitosan films by increasing the degree of cross linking could be because of increasing drug conjugation capacity as the key factor. Also, the morphology of the drug conjugated microspheres acquired by appeared to be little rough and irregular in shape, as evidenced by SEM, probably as a result of the microspheres aggregation. Regarding to the results from this investigation degree of cross-linking and drug conjugation capacity, as two important factors, affect the physicochemical properties drug- and enzyme-biomacromolecules. On the other hand investigation on the properties of biomacromolecules as new carriers is important because of their role in enzyme immobilization studies.

Keywords: Physicochemical Properties, Biomacromolecules, Drug Targeting.

Abstract No.162

How Social Stresses can Alter Microtubular Proteins Structure, Dynamicity and Kinetics

*Ghazaleh Eskandari Sedighi¹, Gholamhossein Riazi*¹,
Mohammadreza Vaez Mahdavi²*

1. Department of Biochemistry, Institute of Biochemistry and Biophysics, University of Tehran, Tehran, IR
 2. Department of Physiology, Shahed University of Medical Sciences, Tehran, IR
- (E-mail: ghazaleheskandari@ibb.ut.ac.ir)

Social stresses are considered as one of the major components of our modern lives. Stressful life events are generally considered as having precipitating effects on the development of human psychopathologies such as anxiety and depression. Previous works and experiments have shown that neuronal plasticity - which is defined as structural adaptation of neurons to functional requirements- can be affected by chronic and acute stress conditions. These structural changes can be