

Abstract No.276

Biophysical Targeting of OmpF Nano-Channel Forming Proteins, Translocating Ofloxacin Antibiotic into E.Coli, Involved In Urinary Tract Infection

*Maryam Rezaei*¹, Hamid Mobasheri²*

1. Institute of biochemistry and biophysics (IBB), IR
2. Lab. Mem. Biophysics, Institute of Biochemistry and Biophysics, University of Tehran, IR
(E-mail: Rezaei@ibb.ut.ac.ir)

OmpF is an ion channel in outer membrane of E.coli that is responsible for translocation of molecules with an exclusion limit of 600Da. These channels are known for the path they form for translocation of hydrophilic molecules, antibiotics, needed to pass through the hydrophobic membrane. An efficient antibiotic is expected to exhibit high affinity for certain target cell and molecules and reach them rapidly. Quinolons and Beta-lactames are among the successful and widely used antibiotics known to cure bacterial infections at present. The antibiotic that enter the bacteria kill them by inhibiting DNA gyrase or the topoisomerase type 2 enzymes as a target, thereby interfering the DNA replication and translocation. In this work, we investigated the passage of Ofloxacin antibiotic through the single OmpF porin of E.coli by means of voltage clamp technique in real time to address the drug trafficking in urinary tract infection disease in human. We also investigated the effect of the passing antibiotic on channel gating, conductance, and voltage sensitivity. The experimental work was further elaborated by theoretical and modeling approaches to address the exact electrostatic effect of antibiotic on the lining group in particular those at eyelet area of the nano-channel that constrict it to a diameter of about 0.4nm. Our results showed that the Ofloxacin is not soluble in lipid phase and did not pass through the membrane. Although the voltage sensitivity of the channel was not changed in the presence of the antibiotics, the frequency of gating increased and fast flickering was caused due to the transient obstruction. The conductance of the channel also remained the same indicating lack of any binding and conformation induction caused by the antibiotic.

Keywords: OmpF Channel, Ofloxacin Antibiotic, Urinary Tract Infection, Translocation.

Abstract No.277

Targeting Mesothelin for Treatment of Cancer

*Kun Wang¹, Vidya Bodempudi¹, Zhengjian Liu¹, Emma Borrego-Diaz¹, Farnaz Yamoutpoor¹, Weihong Pan¹, Arkadiusz Z. Dudek¹, Mojtaba S. Olyaei¹, Tuba Esfandyari², Faris Farassati*³*

1. The University of Kansas Medical Center, Department of Medicine, Division of Gastroenterology, Hepatology and Motility, Molecular Medicine Laboratory, Kansas City, Kansas, US
2. Department of Internal Medicine-Molecular Medicine Laboratory, Divisions of Gastroenterology and Hematology/Oncology, Kansas, US
3. The University of Kansas Medical Center, Department of Medicine, Divisions of Gastroenterology and Hematology/Oncology, Molecular Medicine Laboratory, Kansas City, Kansas, US
(E-mail: ffarassati@kumc.edu)

Mesothelin, a differentiation antigen present in a series of malignancies such as mesothelioma, ovarian, lung and pancreatic cancer, has been studied as a marker for diagnosis and a target for immunotherapy. We, however, were interested in evaluating the effects of direct targeting of Mesothelin on the viability of cancer cells as the first step towards developing a novel therapeutic strategy. We report here that gene specific silencing for Mesothelin by distinct methods (siRNA and miRNA) decreased viability of cancer cells from different origins such as mesothelioma (H2373), ovarian cancer (Skov3 and Ovar-5) and pancreatic cancer (Miapaca2 and Panc-1). Additionally, the invasiveness of cancer cells was also significantly decreased upon such treatment. We then investigated pro-oncogenic signaling characteristics of cells upon mesothelin-silencing which revealed a significant decrease in phospho-ERK1 and PI3K/AKT activity. The molecular mechanism of reduced invasiveness was connected to the reduced expression of β -Catenin, an important marker of EMT (epithelial-mesenchymal transition). Ero1, a protein involved in clearing unfolded proteins and a member of the ER-Stress (endoplasmic reticulum-stress) pathway was also markedly reduced. Furthermore, Mesothelin silencing caused a significant increase in fraction of cancer cells in S-phase. In next step, treatment of ovarian cancer cells (OVca429) with lentivirus expressing anti-mesothelin microRNA resulted in significant loss of viability, invasiveness, and morphological alterations. Therefore, we propose the inhibition of Mesothelin as a potential novel strategy for targeting human malignancies.

Keywords: Mesothelin, Cancer, Immunotherapy, Pro-oncogenic Signaling.