

(E-mail: rr_ss804@yahoo.com)

Phenolic phytochemical compounds are reactive toward free radicals such as reactive oxygen species (ROS). Free radicals and ROS are implicated in protein/DNA damage, cancer and especially accelerated cell aging. Herein, the structural and electronic properties of the four phenolic phytochemical compounds including caffeic acid, ferulic acid, p-coumaric acid and sinapinic acid have been theoretically investigated by performing semi-empirical molecular orbital theory at the level of AM1 quantum chemical method. Structure of the caffeic acid showing features important in defining the classical antioxidant potential of phenolic acids. The results indicated that caffeic acid is more reactive toward free radicals. This is mainly because of the catechol or dihydroxylated ring that having hydroxyl substitutions, enable the compound to scavenge free radicals.

Keywords: Semi-empirical, AM1, Antioxidant, Phenolic Acids, Caffeic Acid.

Abstract No.180

Golden Hamster Monoclonal Antibodies in the Treatment of Metastatic Rectal Cancer in Veterinary Kazeroon, Iran

Mojtaba Norouzi*¹, Mohammad Norouzi², Naser Ghaemi³,
Nour Amirmozafari⁴

1. Department of Microbiology, Islamic Azad University Lahijan, Lahijan, IR
2. Department of Microbiology, Islamic Azad University Rasht, Doctor Veterinary Medicine(DVM), Rasht, IR
3. Department of Biotechnology, University of Tehran, Tehran, IR
4. Department of Microbiology University of Tehran, Tehran, IR
(E-mail: vet_mohamad@yahoo.com)

Monoclonal antibodies(MAB)have found increasing use experimental therapies.great limitation of their use is that they are recognized by the patient as being of foreign origin and an antiglobulin response is provoked.recombinant DNA technology offers the ability to convert these rodent antibodies into a more human form.there are currently several different strategies which can be adopted to generate humanized antibodies resulting in different degrees of humanization can be achieved ranging from chimeric antibodies with a combination of human constant regions with rodent variable regions to fully reshaped antibodies where the variable regions are also humanized. At present the available data on clinical use of chimeric and reshaped antibodies is very limited. The rat IgG2b antibody CAMPATH-1G has

been shown to be both useful as an immunosuppressive antibody as well as in the treatment of lymphoid malignancies. The reshaped version, CAMPATH-1H, was successfully used to clear detectable malignant cells from the blood and bone marrow in two patients with B-cell lymphoma . A more sustained course of the human antibody (126 mg over 30 days and 86 mg over 43 days) was tolerated than had previously been used for the rat antibody.

In the future in veterinary clinic of Sina gilán, it is clear that a majority of monoclonal antibodies produced for therapy will be humanised for the reasons discussed above. As far as improvements in the abilities of these antibodies to interact with human effector mechanisms goes it seems that there is unlikely to be any major differences between chimeric and fully reshaped antibodies.

important about humanised antibodies is whether there are any sequences contained within the variable region frameworks, complimentary determining regions or constant region allotypes which can be processed and presented as T-cell epitopes.

Keywords: Monoclonal Antibodies, Chimeric Antibody, Recombinant DNA Technology, Humanized Antibody, Antiglobulin.

Abstract No.181

Differential Scanning Calorimetry Study of Camel Serum Albumin in Presence and Absence of Fatty Acids

Farzaneh Farivar*¹, Maryam Salami¹, Ali Moosavi-Movahedi¹,
Ali Akbar Saboury¹, Amir Niasari-Naslaji², Mousa Bohlooli¹

1. Institute of Biochemistry and Biophysics, University of Tehran, IR
2. Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Tehran, IR
(E-mail: farivar@ibb.ut.ac.ir)

This study aims to determine the thermal stability of camel albumin and compare it with human and bovine sources. Camel albumin was purified from serum via combination of Cohn method and anion-exchange chromatography. Physiologically albumin binds 0.1-2 fatty acid per mol so activated charcoal treatment was taken to remove them. The thermal denaturation process of camel albumin containing fatty acid (CAF) and free fatty acid camel albumins (CA) in aqueous solution was studied by use of differential scanning calorimetry. The melting temperature of CAF was similar to human serum albumin (80°C) and markedly higher than bovine source (69°C). Removing fatty acids decreased melting temperature of camel albumin, which shows the stabilizing effect of fatty acid. Here fatty acid may be