

descriptors is calculated for each amino acid and they are put together to produce a descriptor data matrix for a set of peptides.

Very recently, in our research group we have been involved in the definition of some new amino acid indices (AAI) for use in QSAM study of peptides. We employed the concept of atom in molecule and used the quantum topological molecular similarity (QTMS) descriptors to suggest some new AAIs. In addition, we employed the Moreau and Broto autocorrelation function on the QTMS descriptors of amino acids to extend the QTMS-based AA indices and generate some other novel indices. In another work, to increase the quality of the QSAR models reported for peptides, we proposed here the application of segmented principal component regression (SPCR) method to define more relevant sources of AA indices. Finally, using of multi-way data analysis methods in QSAM study of peptides will be explained. The suggested methods were validated by analysis of some different peptide data sets of relevant biological activities.

Keywords: Amino acid indices, Peptide, Biological activity, Structure-function.

Abstract No.294

A Possible Role of Peroxidase Activity of "heme-ferritin" Complex in Development of Neurodegenerative Diseases

*Morteza Jafari^{1,3}, Reza Khodarahmi², Samira Ranjbar³,
Mohammad Reza Ashrafi³, Seyyed Mohsen Asghari⁴*

1. Dept. of Biology, Faculty of Science, Guilan University, Guilan, IR
2. Medical Biology Research Center, Departments of Pharmacognosy and Biotechnology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, IR
3. Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, IR
4. Dept. of Biology, Faculty of Sciences, Guilan University, Rasht, IR
(E-mail: mortezajafari1360@gmail.com)

Ferritin is a ubiquitous protein for the sequestration, storage and release of iron in animal and plant cells. It has been previously shown that each ferritin molecule can bind to 15 - 17 heme moieties. Heme and sometimes heme-protein complexes possess peroxidase activities which cause irreversible damages in cells. In the present study, we reported the peroxidase activity of "heme-ferritin" complex using several oxidizable substrates. First, the ferritin was purified from liver homogenate of sheep by ammonium sulphate precipitation and DEAE ion-exchange chromatography. The peroxidase activity of "heme-ferritin" complex was then measured by using H₂O₂ or t-BHP as oxidant

substrates and TMB, L-Dopa or DOPA as a reductant substrates. Results showed that the non-specific peroxidase activity of the "heme-ferritin" system can oxidize the neurotransmitters with high catalytic efficiency. According to the literature, the concentration of ferritin, free heme, and peroxide species increase in neurodegenerative diseases such as AD. There is this possibility that the peroxidase activity of "heme-ferritin" complex play a significant role in development of oxidative damage within involved cells. We will discuss the importance of our observations.

Keywords: AD, Peroxidase Activity, Ferritin, Neurodegenerative Diseases.

Abstract No.295

Interaction of a Novel Photochromic Molecule with Human Serum Albumin Studied by UV-vis Absorption, Fluorescence Spectroscopy, and Docking Calculation

Fereshteh Shaheri¹, Davood Ajloo^{1,3}, Hamzeh Kiyani¹,
Maryam Ghadamghahi¹, Taghi Lashkarbolooki^{2,3}*

1. Department of chemistry, Damghan University, Damghan, IR
2. Department of Biology, Damghan University, Damghan, IR
3. Institute of Biological Science, Damghan University, Damghan, IR
(E-mail: ajloo@du.ac.ir)

Photochromism is the phenomena that of reversible transformation between two forms (open and close form) that have different spectra by photo-irradiation. In this work, the interaction of a new photochromic molecule, 2,2-biphenyl-4-yl-2-methyl-6(4-nitrophenyl)-4-phenyl-1,3-diazobicyclo[3.1.0] hex-3-en, with human serum albumin (HSA) has been investigated by UV-vis absorption, fluorescence spectroscopy and docking calculations. 0.1 μM HSA was used in the presence of 0 to 1.47×10⁻⁵ M photochromic molecule. The interaction of photochromic molecule with HSA causes fluorescence quenching of HSA. The addition of photochrome to a solution of HSA induces quenching of the protein fluorescence and binding constant of interaction was also obtained about 7.5 μM⁻¹. In other investigation the photochromic molecule absorbance was studied by a UV-Vis spectrophotometer model Perkin Elmer. The photochromic reaction was studied upon variation of time irradiation by UV-vis spectrophotometer. Absorption spectra of photochromic molecule and HSA were obtained as the time of irradiation increase from 0 to 70 s. Absorbance intensity was increased due to the close form of molecule that turn in to the open form. Rate constant of reaction was obtained from absorption spectra about 0.026 s⁻¹. In order to estimate the

affinity and binding energy of photochromic molecule to HSA, Autodock 3.0.5 software was used. The negative value of docking energy has revealed interaction of protein with this molecule.

Keywords: Photochromism, Human Serum Albumin, Rate constant.

Abstract No.296

Electrochemical Characterization of Cytochrome C by Using the Electrode Modified with Cadmium Selenium Nanoparticles

Saeid Rezaei-Zarchi¹, Saber Imani², Ali Mohammad Zand³, Yunes Panahi², Amir Nejad Moghaddam², Eisa Tahmasbpour Marzon²

1. Department of Biology, Payame Noor University, Yazd, IR
2. Chemical Injuries Research Center, Baqiyatallah University of Medical Sciences, Tehran, IR
3. Department of Biology, Basic Science Faculty, IHU, Tehran, IR
(E-mail: Imani.Saber@yahoo.com)

Bare surface of the electrode in electrochemical studies of proteins are not suitable. In this case, leads to a decrease in the rate of electron transfer between electrode and protein and protein irreversible adsorption on the electrode surface which is associated with conformational changes and loss of protein activity. Therefore, you must provide the necessary groups on the electrode surface for active communication with the macromolecules. Molecules could provide the group called facilitator. In this paper, was diagnosed with cytochrome c with modified glass carbon electrode surface by cadmium selenium nanoparticles. This detection can be used in Biosensors designed for measurement of hydrogen peroxide. X-ray Diffraction (XRD) results showed size nanoparticles synthesized with electrochemical method are 57 nm. Alpha

Absorption spectra revealed the size limit. Advantage of electrochemical methods for synthesis of nanoparticles of cadmium selenium is an easy, cheap and low cost. Biosensors with cytochrome c stabilized on this electrode shows high sensitivity and linear range from 15 to 700 micro M for determination of hydrogen peroxide.

Keywords: Cytochrome C, Cadmium Selenium Nanoparticles, Electrochemical Characterization.
