

Molecular Analysis of *Vibrio Cholerae* El Tor Variants Isolated from Outbreaks During 2005 in Iran by REP-PCR and ERIC-PCR

Ali Sohrab Nejad*¹; Mostafa Akbari Qomi²; Abbas Maleki³; Sobhan Ghafurian³; Norkhoda Sadeghifard³

1-Department of Public Health, University of Medical Sciences, Ilam, Iran

2-Department of Microbiology, Faculty of Medicine, Ilam University of Medical Sciences, Ilam, Iran

3-Clinical Microbiology Research Center, Ilam University of Medical Sciences, Ilam, Iran

sohrabi1354@yahoo.com

Background & Objectives: In the present study, we analyzed the PCR patterns of *Vibrio cholerae* El Tor variants isolated from outbreaks during 2005 in Iran by REP-PCR and ERIC-PCR.

Methods: Thirty nine El Tor variants were obtained from outbreaks during 2005 in Iran. After detection of isolates by biochemical methods, and serotyping, chromosomal DNA was extracted by standard phenol/chloroform methods. REP-PCR was carried out using Rep 1R- I and Rep 2-I primers, also ERIC-PCR was done by using ERIC -1R and ERIC - 2. The PCR product was run and visualized in 1.5% agarose gels stained with ethidium bromide.

Results: Thirty nine El Tor variants were analyzed and PCR Product sizes by REP-PCR methods were between 250-3500 bp and 5 different PCR patterns were detected. The most common pattern by REP-PCR was observed among 26 (66.7%) isolates. By ERIC-PCR 7 different patterns were detected.

Conclusion: Results of this study showed that most of strains have similar pattern by REP-PCR and ERIC-PCR. It is assumed that these strains have the same origin.

Keywords: *Vibrio cholerae*; Molecular Typing, Iran

